No.



9200253

KHEDER CONTRIBED STRANGES OF FRANCES RECEA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

The Thio State University,

Thio Agricultural Research and Bevelopment Center

Thereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to extude others from selling the variety, or offering it for sale, or reproducing it, importing it, or exporting it, or using it in producing a hybrid or different ty therefrom, to the extent provided by the Plant Variety Protection Act. United States seed of this variety (1) shall be sold by variety name only as of certified seed and (2) shall conform to the number of generations the owner of the rights. (84 Stat. 1542, as amended, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Freedom'

In Lestimony Watercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this 31st day of May in

this 31st day of May in the year of our Lord one thousand nine hundred and ninety-bour.

Kennett Hevan

Commissioner

Plant Variety Protection Office

Agricultural Marketing Service

Citu EST Socrotary of Agriculture Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Expires 1/31/91

U.S. DEPARTMENT OF AGRICULTURAL MARK	AGRICULTURE		Application is required in order to					
APPLICATION FOR PLANT VARIET	TY PROTECTIO	N CERTIFICATE	determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).					
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNATION O	R 3. VARIETY NAME					
The Ohio State University, Ohio Agri	cultural	EXPERIMENTAL NO.						
Research and Development Center		ОН413	Freedom					
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area code)	FOR OFFICIAL USE ONLY					
	i e		PVPO NUMBER					
1680 Madison Avenue			0000057					
Wooster, OH 44691		216-263-3886	9200253					
			F Date					
	,		1 August 25, 1992					
6. GENUS AND SPECIES NAME	7. FAMILY NAME (Bota	nical)	Yime N					
Triticum aestivum, L	Graminae		G 3:50 A.M. PP.M.					
8. CROP KIND NAME (Common Name)	P KIND NAME (Common Name) 9. DATE OF DETERMINATION							
	E : 2150. º							
Soft Red Winter Wheat	S Daye							
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGA	7 1 August 24, 1992							
Agricultural Experiment Station	C Certificate Fee:							
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	- [:250. ºº							
	V Date							
			5 april 18, 1994					
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO	SERVE IN THIS APPLICAT	TION AND RECEIVE ALL PAPERS	,					
Dr. H. N. Lafever	•	•						
Agronomy Dept.	_							
Ohio Agricultural Research & Develop								
1680 Madison Ave., Wooster, OH 4469		PHONE (Include area	code): 216-263-3886					
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (For	low INSTRUCTIONS on revo	975e)						
a. X Exhibit A, Origin and Breeding History of the Variety.								
b. X Exhibit B. Novelty Statement.			4					
c. X Exhibit C, Objective Description of Variety.								
d. X Exhibit D, Additional Description of Variety.								
e X Exhibit E, Statement of the Basis of Applicant's Ownersh		,						
t. X Seed Sample (2,500 viable untreated seeds). Date Seed			losed)					
g. X Filing and Examination Fee (\$2,150) made payable to "1								
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SC Protection Act.) ▼ ▼ ▼ ▼ ▼ ▼ ▼ ▼ ▼ ▼			(See section 83(a) of the Plant Variety					
X YES (If "YES," answer items 16 and 17 be		NO," skip to item 18 below)						
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS NUMBER OF GENERATIONS?	TO 17, IF "YES" 1	TO ITEM 16, WHICH CLASSES OF PRO	DUCTION BEYOND BREEDER SEED?					
X YES NO.	¦ ⊠ Fo	UNDATION X REG	ISTERED X CERTIFIED					
			-					
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VA	RIETY IN THE U.S.?							
YES (If "YES." through Plant Variety Protection Act	Patent Act. Give da	ate: .)						
X NO		-						
10 HACTHE VARIETY DECK DESTANCE LIGHT OFFICER CO.								
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR M	IARKETED IN THE U.S. OR	OTHER COUNTRIES?						
YES (II "YES," give names of countries and dates)								
🛛 NO (Some seed under our control	was distribu	ted to other state	seedstock organizations					
for increase (first on 8/27								
20. The applicant(s) declare(s) that a viable sample of basic se request in accordance with such regulations as may be appl	eds of this variety wil	l be furnished with the applica	tion and will be replenished upon					
the contract of the contract o	•		and the state of the second state of the state of					
The undersigned applicant(s) is (are) the owner(s) of this uniform, and stable as required in section 41, and is entitle	d to protection under t	hover plant variety, and belie he provisions of section 42 of th	e Plant Variety Protection Act.					
Applicant(s) is (are) informed that false representation here								
SIGNATURE OF APPLICANT [Owner(s)]			Love					
11 1 1	CAPACITY OR)	DATE					
Howard II. Labour	Boon A	i QA	6/30/00					
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR	TITLE	DATE					
C	A A	mice .	DATE					
James H / Now	(let.	1.0.0	1.71,5191					

FORM CSSD-470 (5-89) Edition of FORM LS-470, 3-86, is obsolete

Exhibit A

Origin and Breeding History of the Variety

- 1. Freedom (previously known as OH413) originated at The Ohio State University, Ohio Agricultural Research and Development Center from the cross: GR876/OH217, $(OH217=Logan^*3/3/Va63-52-12/Logan//Blueboy-F_2)$. The cross was made in 1980. Freedom was first selected as an F_3 plant in 1983. Fifty heads were reselected in F_7 in 1987 and progeny rows were examined for uniformity and yield through 1990. Its earlier experimental designation was 11880-10.
- 2. Breeder seed of Freedom consists of the progeny of 19 different F_7 plants bulked after harvest in the F_{10} generation in 1990 after the progeny appeared identical and true breeding in separate plots in 1988 through 1990. Seed was bulked after harvest in 1990 and further increased as breeder seed in 1990-91. Foundation generation seed is under production in 1991-92.
- 3. Freedom appears to be very uniform and homozygous as observed in the field in 1990, 1991, and 1992. This uniformity is to be expected with our purification and increase system.
- 4. Freedom appears to be very stable and true breeding as evidenced by various agronomic and pathological examinations conducted on the F_{10} - F_{12} generations in the special purification and increase nurseries.
- 5. Variants observed during the development of this variety were very few in number and of various, non recurring types, typical of breeding programs involving self-pollinated crops
- 6. This variety was selected for its high yielding ability, excellent straw strength, and excellent disease resistance. The variety possesses the 2BL/1RS translocation, the same translocation present in its parent, GR876.

Exhibit B

Novelty Statement and Botanical Description of the Variety

Freedom is an awnless cultivar, but like most soft red winter wheats, it possesses apical awnlets. Freedom appears most similar to Cardinal under normal field conditions in the Midwest, possessing very similar medium green foliage color and head color. Freedom and Cardinal possess apical awnlets of similar length. Spikelets of Freedom are slightly smaller than those of Cardinal and the heads are slightly denser for Freedom and slightly smaller. Freedom tends to possess 2-3 extra spikelets per head compared to Cardinal.

Freedom is a moderately early to mid-season cultivar and is moderately short, averaging 5 cm shorter and 2 days earlier than Cardinal in 18 tests from 1988-1990. The cultivar has exhibited very good straw strength in Ohio tests, being approximately equal to Cardinal and Dynasty. Winterhardiness of Freedom is excellent, equalling that of the best currently grown cultivars in Ohio tests.

Test weight of Freedom is relatively low, averaging approximately .5 lb/bu below that of Excel and slightly more than 1 lb/bu below that of Cardinal.

The USDA Soft Wheat Quality Laboratory, Wooster, Ohio in evaluations of samples of Freedom over a six-year period found Freedom to possess average milling quality and slightly below average baking quality.

Disease resistance of Freedom is excellent; the cultivar having shown no mildew ($Erysiphe\ graminis$) in trials over the past seven years. USDA Cereal Rut Laboratory tests involving several different races of $Puccinia\ recondita$ and $Puccinia\ graminis$ show Freedom to possess excellent resistance to these two pathogens. Ohio field tests have also verified these results. Freedom has also shown very good resistance to wheat spindle streak mosaic virus (WSSM). Freedom possesses resistance to races GP, A, C, E, and F of Hessian fly ($Mayetoila\ destructor$, Say) imparted by the H_3 gene.

Freedom also possesses good to excellent tolerance to acid soil being similar to Cardinal in this regard.

ź

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE COMMODITIES SCIENTIFIC SUPPORT DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

INSTRUCTIONS: See Reverse. WHEAT (TRITICUM SPP.)
HAME OF APPLICANTIA The Ohio State Universit	
Agricultural Research and Development Cent	9200253
	VARIETY NAME OR TEMPORARY
1680 Madison Ave.	DESIGNATION
Wooster, OH 44691	Freedom
Place the appropriate number that describes the varietal chara	cter of this variety in the boxes below.
Place a zero in first box (c.s. 0 8 9 or 0 9) when numb	er is either 99 or less or 9 or less.
1. KIND: 1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT	S = POLISH 6 = POULARD 7 = CLUB
2. TYPE,	
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 = SOFT 3 = OTHER (Specify) 2 = HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specily)	
1. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
2 2 5 FIRST FLOWERING	2 3 1 LAST FLOWERING
4. MATURITY (50% Flowering):	•
NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
3 NO. OF DAYS LATER THAN	1 4 = LEMHI 5 = NUGAINES 6 = LEEDS
5. PLANT HEIGHT (From soil level to top of head):	
9 I CM. HIGH	
CM. TALLER THAN	
	1 = ARTHUR 2 = SCOUT 3 = CHRIS
3 CM. SHORTER THAN	1 4= LEMHI S = NUGAINES 6 = LEEDS
& PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
2 I TYELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT	2 Vaxy bloom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Intermodes: 1 = HOLLOW 2 = SOLID
4 NO. OF NODES (Originating from node above ground)	2 2 CM INTERNODE LENGTH BETWEEN FLAG LEAF
*. AURICLES:	
Anthocyanin: 1 = ABSENT 2'= PRESENT	2 Hairiness: = ABSENT 2 = PRESENT
10. LEAFI	
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
1 Hairs of first leaf sheath: = ABSENT 2 = PRESENT	2 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 2 MM. LEAF WIDTH (First loof below flag 1000	2 9 CM. LEAF LENGTH (First loof below fine loof):
FORM LMGS 470-6 (6-82) (Formerly Form LPGS 470-6 (3-79), which	may be used)

11. HEADI 2 Density: I = LAX	x ? = DENSE	Shape: 1 = TAPE 1 = OTHE	RING 2 = STRAP 3 = CLAVATE
2 Awnedness: 1 = 4	AWNLESS 2 = APICALLY AWNLETED	J = AWNLETED 4 = AWN	εο
2 Color at maturity:		4 = RED HER (\$p+ci(y):	·
8 3 CM. LENGTH		1 2 мм. wioth	
[0:12]			·
: 3 I _	RITY: RT (CA. 7 mm.) Z = MEDIUM (CA. 8 mm.) RT (CA. 9 mm.)	3 = WIDE (OW (CA. 3 mim.) 2 = MEDIUM (CA. 3.5 mim.) CA. 4 mm.)
Shoulder I = WAN 2-4 shape: 4 = SQU	_	Besk: I = OBTUS	E 2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLO	OR:	14. SEEDLING ANTHOC	Ү АНН:
1 1 = WHITE 2 =	RED 3=PURPLE	1 I = ABSENT	2 = PRESENT (occas. trace)
IS. JUVENILE PLANT O	ROWTH HABIT:		•
1 = PROSTRATE	2 = SEMI-ERECT 3 = ERE	ECT	
16. SEED:			
1 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	1 Check: 1 = ROUNG	DED 2 = ANGULAR
1 Brush. 1 = SHORT	2 = MEDIUM 3 = LONG	Brush: I = NOT C	OLLARED 2 = COLLARED
NA (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK	WN .	
3 Color: 1 = WHITE	2 = AMBER 3 = REO 4 = PURPLE	5 = OTHER (Specily)	
7 MM. LENGTH	4 MM. WIDTH	3 5 GM. PER 1000	SEEDS
17. SEED CREASE:			
1 Vidth: 1 = 602 OR	LESS OF KERNEL 'WINOKA'	1 1 1	R LESS OF KERNEL 'SCOUT'
	ESS OF KERNEL 'CHRIS'		R LESS OF KERNEL "CHRIS" R LESS OF KERNEL "LEMH!"
	AS WIDE AS KERNEL "LEMHI"	3 - 30 1 07	T C C S S T T C T T T T T T T T T T T T
stem Rust See	LEAF RUST See (Reces) attached	O STRIPE RUST	O LOOSE SMUT
2 POWDERY MILDEN	O BUNT	OTHER (Specify)	
9. INSECT: (0 = Hot Took	ed, 1 = Susceptible, 2 = Resistant)		
SAWFLY	0 APHID (Bydv.)	O GREEN BUG	O GEREAL LEAF BEETLE
OTHER (Specify)	HESSIAN FLY	0 GP 2 A	1 B 2 c
	RACES:	1 o 1 E	2
	ETY MOST CLOSELY RESEMBLES THAT S	UBMITTED:	RECEIVED V
CHARACTER	NAME OF VARIETY	CHARACTER	HANE OF VARIETY
Plant tillering	Cardinal	Seed size	GR 876
Leaf size	GR 876	Seed shape	GR 876 2 4 1992
Leaf color	Cardinal	Coleoptile elongation	GR \876
Leaf carriage	Cardinal	Seedling pigmentation	Cardinal Protection Ofc
	INSTRU	J 1 10.75	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

(a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Theat Varieties Grown in the United States. Technical Bulletin 1278, United States Department of Agriculture.

⁽b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysis. (See attachments)

The Ohio State University Ohio Agricultural Research and Development Center Wooster, Ohio

RELEASE OF FREEDOM SOFT RED WINTER WHEAT

The Ohio State University, Ohio Agricultural Research and Development Center announces the release of 'Freedom' soft red winter wheat, a new, productive cultivar for Ohio and surrounding states.

Freedom (previously designated OH413) resulted from the cross: GR876/OH217 (OH217 - $Logan^{\dagger}3/3/Va$ 63-52-12/ $Logan/Blueboy-F_2$). First selected in 1983 as an F_3 plant. Fifty heads were reselected in F_7 in 1987 and progeny rows were examined for uniformity and/or yield through 1990. Breeder seed consists of progeny of 19 different F_7 plants bulked after harvest in the F_{10} generation in 1990.

Freedom was first tested in Advanced Ohio Trials in 1986. At the end of the 1990 season it had been tested in 22 state-wide, drilled plot trials over a 5-year period. Freedom was also an entry in the Uniform Eastern Soft Red Winter Nursery in 1989 and 1990.

Freedom is a beardless, white chaffed cultivar. It is a medium short cultivar of medium maturity. Straw strength of Freedom appears equal to Cardinal and Dynasty. Winterhardiness of Freedom is excellent, being equal to Becker, Dynasty and Excel. Yields of Freedom have exceeded Becker and Cardinal by several bushels, but averaged only 0.6 and 0.7 bu/a above Dynasty and Excel, respectively, in 18 tests over a 3-year period. (See Tables 1-4 for additional data.)

The USDA Soft Wheat Quality Laboratory at Wooster, has reported Freedom to possess acceptable milling and baking quality based on tests over 6 years. Quality was considered to be average.

Freedom possesses excellent resistance to both leaf rust (Puccinia recondita) and powdery mildew (Erisiphe graminis) in field evaluations in Ohio. It is also highly resistant to wheat spindle streak mosaic virus (WSSM). It also appears resistant to all selected isolates of stem rust (Puccinia graminis) according to published 1990 Uniform Eastern Soft Red Winter Nursery results conducted by the USDA-ARS Cereal Rust Laboratory. In 1989 and 1990 Freedom exhibited few septoria leaf or glume blotch symptoms under field conditions in Ohio even though these diseases were severe both years in terms of overall nursery infections. Tests conducted by the USDA-ARS Hessian Fly Lab, Lafayette, IN indicate Freedom possesses only the H3 gene for Hessian fly resistance.

Freedom was so named in honor of the troops who participated in Operation Desert Storm as well as seeming appropriate in view of its relative freedom from disease; presumed to be largely due to its possessing the 1B/IR translocation from rye.

Application for Plant Variety Protection will be made for Freedom. The classes Foundation, Registered, and Certified will be allowed beyond Breeder seed. Breeder seed of Freedom will be maintained by The Ohio State University, Ohio Agricultural Research and Development Center, Wooster, OH, 44691.

Release of information to the general public regarding the name, release, performance or description of Freedom by those participating in the release may be made after August 1, 1991.

Freedom will be available as Breeder seed to other states for fall, 1991 seeding.

States interested in the increase and release of this new cultivar must notify Dr. H. N. Lafever, Agronomy Department, Ohio Agricultural Research and Development Center, Wooster, OH, 44691 in writing by August 15, 1991 indicating the quantity of Breeder seed desired. Pricing will be appropriate for this class of seed.

A royalty fee of \$ 0.25 per bushel must be assessed and collected on the Registered and Certified class of seed of Freedom by states participating in release of this cultivar. Research and development fees collected by another state will be shared equally, with 50% retained by that state, if allowed by the Director, to be used for agronomic crop breeding research and development. The other 50% is to be returned to the Ohio Agricultural Research and Development Center. It is recommended that an additional \$.10 per bushel administrative fee be levied by participating states to cover the cost of collection, advertising, and protection of the cultivar in the state of collection. Contractual arrangements will be completed at a later date with those states participating in the release of Freedom.

James H. Brown, Acting Director The Ohio State University

Ohio Agricultural Research and

Development Center

July 18, 1991

Table 1. Comparison of yields of Freedom with currently grown cultivars in drilled plot trials, by years, Ohio.

							Summary	
Cultivar	1986 1 test	1987 3 tests	1988 6 tests	1989 6 tests	1990 6 tests	Avg. 21 tests	Avg. 18 tests	Avg. 15 tests
			¥ *****		1.10.20.04	****		· ·
Freedom	70.2	76.0	69.6	64.9	74.7	70.6	69.7	70.2
Becker	49.1	64.8	67.4	57.4	67.4	64.2	64.1	63.2
Caldwell	49.5	62.4	67.9	54.2				61.5
Cardinal		68.5	60.3	60.4	72.1	64.9	64.3	63.1
Dynasty	· 		71.5	66.5	69.4		69.1	
Excel			71.3	65.5	70.1		69.0	
5% L.S.D.	4.5	5.7- 6.9	3.0- N.S.	4.3- 6.7	3.7- 5.2			

Table 2. Comparative Agronomic performance of Freedom and currently grown cultivars in drilled plot trials, Ohio (average of 18 tests, 1988-90).

		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~		
inter Survival (%)	P1. Ht. (in.)	Date Headed (May)	Lodging (%)	Test Wt. (lb/bu)
96	36	25	8	54.9
96	33	26	3	55.9
91	38	27	7	56.2
96	38	25	8	57.7
96	36	26	2	55.5
	96	96 36	96 36 26	96 36 26 2

Comparative performance of Freedom and currently grown cultivars in Table 3. miscellaneous Ohio tests.

	H.F.	% Mildew ¹	WSSM ²	Leaf Rust	Al Tol		Quality	<u>- 6 yrs</u>
Cultivar	Res. (Races)	12 tests 5 yrs	4 tests 1 yr	4 tests 4 yrs	Yield ³ 1 yr	Score ⁴ 5 yrs	Milling	Baking
Freedom	A,C,F	0	2	0		4.8	. <b>C-</b>	D
Becker	A,C,F	82	1	3MR	100	4.0	С	В-
Caldwell	A,B,E	29	4	OVR	100	6.3		
Cardinal	A,C,F	33	2	1VR	100	4.8	A ·	В
Dynasty	None	13	1	2VR	43	6.5	В-	D
Excel	A,B,C,E,F	13	1	4VR	97	7.2	B-	C-

¹ Years or locations with no mildew present omitted from calculations.
2 0 - none to 9 - severe.
3 Yield as a % of Seneca.

 $^{^4}$  0 = very tolerant to 9 = sensitive.

Summary of Regional Nursery data involving Freedom and currently grown cultivars in Ohio tests (4 tests).

						Ave	744	1000			:	
Cultivar	196	Yie	Yield (bu/a)	/a)	Avra	Date Date	Avg. Ht.	Avg. Lodging S	Avg. Survival	Avg. Mildew	H.F. Races	Test Wt.
	OARDC	N.W.	OARDC N.W. OARDC N.W.	N.W.		(May)	(in.)	(in.) (%)	(%)	(%)	ગ પ્ર	(1b/bu)
												and the first firs
Cardinal	74.1	73.4 52.0	52.0	53.5	53.5 63.3	28	39	20	06	32	R-S-S	53.2
Excel	71.8	9.07	70.6 51.5	6.49	64.7	27	36	17	92	16	R-R-S	52.2
Freedom	75.2	75.9 53.7	53.7	6.69	68.7	. 27	37	11	89	0	R-S-S	52.2
Wakefield	80.4	75.2 58.7	58.7	68.1	9.07	29	39	22	98	0	S-S-S	53.6
Caldwell	9.79	61.5 38.6	38.6	50.2	54.5	26	36	25	89	21	R-R-S	52.7
5% L.S.D.	5.6	0.9	6.0 9.4	11.4								

Seedling reaction of entries of the 1990 Uniform Eastern Soft Red Winter Wheat Performance Nursery to selected isolates of <u>Puccinia graminis</u> f. sp. <u>tritici</u>. (by D.V. McVey, USDA-ARS, Cereal Rust Laboratory, U. of MN., St. Paul, MN.)

Maria de la companya del companya de la companya de la companya del companya de la companya de l	<del></del>	React	ion Pr	oduced	by Is	olates		<u> </u>
	68-	72-	69-	72-	72-	72-	74-	
	41-	00-	21-	25-	00-	4 -	21-	
	73A	1370C	399	639C	53A	lA	1409A	
Name or	<u>HNLO</u>	QFBS	QSHS	RKQS		TNMH		Spec.
No. Sel. No.	17A	15			32	153		Sr Gene
1 Knox 62	s s	S	S	S	S	S	S	NONE
2 Cardinal	S	S	S	s	S	S	S	NONE
3 Caldwell	0 .	2=	S	S	٥	0	2=	9a,17
4 OH 236	0	2=	2=	S	0;	0	2= S 2-	7b,17
5 AR 26415	2=	2=	S	211	2-	2 S,0;	2-	<b>+</b>
6 OH 394	0	S	0,5	S	S	S,0:	S	5,seg 17,36?
7 KY 83-38	S	S	s ·	S	S	s	S	иоиЕ
8 MO 10501	2=	2=	S	S	S	2=	2-	9a
9 ABI 85-81	Ō	<u>1</u> N	ō	;1C			;1CN	6,17,36
10 MO 11769	Š	s	Š	S	s	; S	S	NONE
11 MO 11785	Õ	;	s	;,s	0	0	2	
12 OH 413 (Freedom)	Ö	ó;	0	0,3	0	0	0	10,17,÷
13 VA 85-52-24	0	0,	0	0				+
14 VA 85-52-34	; lN	_		_	0	0	0	+
15 IL 84-3010		; lN	S	;lN	īN	;1-N	;1CN	10,15,17
16 IL 84-3511	S,;1		s	s	Ş	S,;	S	seg 17
	;l=N	23CN		S	S	;1N,S		seg 17
17 PA 8457-1	2=	2=	S 2	S	S	2-	2-	5,9a
18 WI X1625-1-1	, 2-	2=	2	S	ឆននេច	s s	S	+
19 MD 75266-46	0,5	S	s o	S	S	S	s s o	seg 5
20 C 86-33	0	0	0	0		0		4.
21 CL 860426 ·	0	ı s s	0	;1	0	0	0 S S S	+
22 MD Blend	s	S	S	S	S	S	· S	NONE
23 Susquehanna	0,5	s	s s	s s	S	s s	S	seg 5
24 T 84-774	0	s,;	S	S	S	;	S	+
25 T 84-331	0	o o	23	;1CN		Ö	n ′	6,10
26 79410D1-3-3-5-2-	1 0	s	0	s	o'	ŏ	2=	17,36,+
27 813811-16-5-50	0 .	s s	S	Š	ŏ ·	Ŏ	S	5,17
28 7942H1-20-8	0	s	S	s	0;,5	;,s		5,seg 17
29 ABI 86-91	Ō	2-11	s	ō	;1-N	s	s s	5,10
30 ABI 86-55	ō	0	Š	ŏ	;1-N	S	S	5,10
31 PSR-W71	Ö	š			0	0		5,10
32 PSR-W84	Ŏ	s	s s	s s s	ŏ	0	S	17
33 PSR-W89	ō	s	S	2	0	0	S	17
34 IL 84-4046	2	2	S	S			S	17
35 WI X1625-1-3	2	2-	ა 2≔	<u>ی</u>	S	2-	2-	9b
36 WI X1923-1	ő	2		2	2-	s	S	+
37 VS 30409	Ö	s S	23	S	S	2-	2	5,9a
38 VS 30557	s	S	S	S	0	0	S	17
39 C 86-24			S	S	S	S	S	NONE
40 LB 63	;1-	·	S	S	-	;1-CN		+
41 AR 84-31-5	0	2=	2-	2≔	0	0	2=	17,24
MC 04-21-2	. 0	S	S	S	0	<b>O</b>	S	17

					Reac	tions	produce	ed by	NA rac	е	<u></u>		
: <del>=</del>	Cultivar or line	TLGG ¹	MBGL	TBGL	CBGL	CBGB	DBBB	LBBQ	MFBL	MDBL	PBRG	PLML	Postulated Lr genes
	Knox 62	3	3	3	3	3	3	3	3.	3	3	3	0
	Cardinal	;	3	1c2	1c;	;	;lc	3	;lc-:	3 3;-10		21c	+ '
	Caldwell	3;	3	3	3	3	;lc	3	3	3	lc:	1c2	+
	OH 286	3	3	3	3	3	;1c	3	3	3	3	3	+
	AR 26415	3	3	3	3	3	;	; 2	; 1	; 1	3	lcl	- 11
	OH 394	;	;lc	;lc	;1c	;	; 1	;	;	;1c	3	3	+
	KY 83-38	3;1c	3	3	3	3;	3;	3	3	3	3	3	+
	MO 10501	3	3	3	;		;ì	;	3	3	3	3	1 ,
	ABI 85-81	•	1c;	;lc1	:	•	:		1c2	;lc	21	1c;2	2
	MO 11769	3	3	3	, 1c	:	,1c	;1c	:1c1	;1c1	3	21;	1,11
	MO 11785	3-1c2	3-:	;lc-3	•	3-;	3;	3;	3	•	1c3;	31c	+
	OH 413 (Freedom)		• •	,	•	:	•	;1c	3:		:	:	+
	VA 8552-24	•	3;	3;	3;	•	•	,		:	:	;1	10,11
	VA 85-52-34		3	3	3,	•		3	3	ź	3	á ·	10
	IL 84-3010	, 2	3	. 3		: - 3	: - 3	ৰ	3	3.	3	3	+
	IL 84-3511	- 3	3	3	•	:3	: - 3	3	จั	3	3	3-;	+
	PA 8457-1	3			•		, - J		•	•		3	9
	VI X1625-1-1	3;	,	,	3	, .	,	,	,	3	3	3	ž
	MD 75266-46	3, 3	3		3	ر ع	3	,	3	3	3	3	Ō
	MD 75266-46				_		-			;lc		•	• •
	CL 860426	3	3	3;	;c ;lc	;c ;1c2	c		•	, 16	, 2	•	1,3,11
-	_	:-3		٥,		; 1C2	1			3	3	, a	4
	MD Blend	;-3	3-; 3-;	-	3-;	į	c	•	,	3-3;	3-3;	3	· -
	Susquehanna	į		3	3;		.1. 2		3-;	;2c	3-2,	3	
	T 84-774	į	;1c	;lcl	;lc	;1c	;1c-3	110	;1c	, 26			2a,11
	T 84-331	3	į	3	;	;	;	3 7	Ş	i	,	•	7 <b>4</b> ,11
	79410D1-3-3-5	; - 2	3	3	3	; 3	;	11c;	2	1+		, 1	T 1.
	813811-16-5-50	;1c3	3	31c	;lc	;1c	;1c2	;_	3;	1c;-3		lc;	+
	79424H1-20-8	;lc-3	3;	;	;1c	; 3	;1c	;1c	;_	2;	1c;-3		
	ABI 86-91	3	3	-	3	3	<u>;</u>	;1c	;lc	į	3;	;lc	11
	ABI 86-55	3;	3	3	3	3	3-;	; - 3	;	3;	3	;1c-3	
	PSR-W71	3-;1c	<u>,3</u>	3	;	;	;	3	;-3	3	3	3	1
	PSR-W84	;lc	3	3	lc;	31c;	;1c	;	;	;	į	;	+
	PSR-W89	3	3	3	lc;	;	;	; 2	;	;	3	;	1,11
	IL 84-4046	3	;	3	;	;	;	;	;	;	; 1	; - 3	1,2a
	₹I X1625-1-3	1c2	3	3	3;	3;	;	•	3;	3	3	3	+
	₹I X1923-1	3	3	3	3	3	;	;	3	3	3	3	3
	/S 30409	3	3	3-;	3	3	;1c	1+	3	3	3	3	+
	/S 30557	;1c3	3	3	3	3;	31c	3;	3	3	2;-3	3;	+
	86-24	;	•	;	;	;	-		•	;	;	-	• •
	.B 63	;	;	<b>;</b> ·	;	;	;	1c2	; - 3	;	;	;	+
A	R 84-31-5	· <b>;</b>	;1c	;	;	;	;	;	;	;	;	;	
													_

Single <u>Lr</u> genes tested - 1,2a,2c,3,3ka,9,10,11,16,17,18,24,26,30.

Virulence formula:

5 7 8

0

TLGG - Lr1, 2a, 2c, 3, 9, 11, 18

MBGL - Lr1, 3, 10, 11

TBGL - Lr1, 2a, 2c, 3, 10, 11

CBGL - <u>Lr</u>3,10,11

CBGB - Lr3,11

DBBB  $\rightarrow Lr2c$ 

 $LBBQ - \underline{Lr}1,10,18$ 

MFBL = Lr1,3,10,24,26

MDBL - Lr1, 3, 10, 24

PBRG - Lr1, 2c, 3, 3ka, 11, 18, 30

- PLML - <u>Lr</u>1,2c,3,3ka,9,10,30

### Exhibit D

### Additional Description of the Variety

Heads of Freedom are best described as dense (Item 11, Exhibit C), however, they are only moderately dense compared to many varieties described as dense.

Freedom is apically awnletted with tip awns measuring 20-35 mm in length.

Auricles of Freedom are typically free of anthocyanin, however, in some seasons anthocyanin in auricles will occur, especially at row ends.

### Exhibit E

### Statement of the Basis of Applicant's Ownership

The originating cross, early line increase and evaluation, selection, reselection, testing, purification, and final multiplication were all performed by the applicant breeder (Dr. H. N. Lafever) or his technical assistants on the property of The Ohio State University, Ohio Agricultural Research and Development Center utilizing funds provided for such research. The variety is intended for release as a public variety in the United States.